

ACRP

Final Report – Activity Report

Program control:

Climate and Energy Fund

Program management:

Kommunalkredit Public Consulting GmbH (KPC)

1 Project Data

Short title	CC-IMPATY	
Full title	Climate Change impact on humidity	
Project number	B286211	
Program/Program line	ACRP 5 th Call for Proposals	
Applicant	Zentralanstalt für Meteorologie und Geodynamik Mag. Barbara Chimani, successor of Dr. Ingeborg Auer since February 2014	
Project partners	University of Bonn	
Project start and duration	Project start: 18.02.2013	Duration: 30 months
Reporting period	from 18.02.2013 to 17.08.2015	
Synopsis: The characteristics of the time series of relative humidity (daily averages) and possible breaks have been investigated and according to these results, different benchmark datasets have been created, representing real characteristics of inhomogeneous time series, but with known inhomogeneities and break locations. These datasets have been homogenised during a blind test by 17 homogenisation methods (different combinations of break detection and break correction and different configurations). Most of the methods underestimate the number of breaks in the time series. For the correction of the time series two of the methods (MASH and ACMANT) show similar ability. ACMANT has been used to		

ACRP — Calls for Proposals

homogenise Austrian station data of relative humidity. The homogenisation showed a strong impact on the trend of relative humidity. The data was used for different analyses with stakeholders concerning gradients and pollutants as well as for climatological analyses.

2 Technical /Scientific Description of the Project

2.1 Project abstract (max. 2 pages)

Daily homogenisation is of more and more importance for climatological studies. During the last years, the abilities of different methods to homogenise daily data of temperature and precipitation have been developed and tested. Nevertheless, other parameters such as humidity can also be important, and therefore different methods have been tested in this project for their ability to homogenise this parameter. From all the possible humidity parameters, relative humidity (daily averages) was chosen, as this parameter is the most widely used by the stakeholders of the project. Humidity is not only of climatological importance, but also has an influence on building materials, different plants and the ability to produce artificial snow.

The four different homogenisation methods used have been used in the past in the COST-Action ES0601 and tested for temperature and precipitation. As most of the methods can be used with different settings, 17 different results for the homogenisation tests have been produced. In order to get an objective estimate of the ability of the different methods, a surrogate dataset containing different difficulties was created. This dataset reproduced the statistical characteristics of relative humidity and possible breaks. This statistics resulted from studies of parallel measurements and the available metadata of the Austrian stations (including number of relocations and instrumentations changes). The ability of break detection and correction has been analysed. Afterwards the best of these methods has been implemented to homogenise real Austrian station data.

Furthermore, the relative humidity data, together with the homogenised temperature data of the ACRP HOM-START project (K09AC0K00025) was used for different analyses done together with the stakeholders and in climatological studies showing the effect of the homogenisation.

Due to the small break signal in the relative humidity time series, the detection of these breaks is challenging for all the methods. The highest efficiency in detecting breaks was found using ACMANT and HOMOP. For the assessment of the quality of the correction of time series the parameters RMSE, variance and trend deviations have been used as criteria. ACMANT and MASH showed the best correction results in the realistic case of the surrogate data.

Due to its ability to accurately detect and correct, ACMANT was chosen for homogenising the Austrian relative humidity data. A homogenisation of 34 stations was possible. Without taking individual station inspection into account, about half of the breaks found in the time series could be confirmed by metadata.

The analyses of the homogenised data showed that the trend in most of the time series grew stronger, having a negative slope during the measuring period of the individual time series,

ACRP – Calls for Proposals

starting in 1948 in the earliest for these homogenisations. This is in agreement with the dataset homogenised in HOM-START, where the same maximal starting year was used and the same pool of stations was homogenised. The homogenisation led to higher medians of relative humidity during all months of the year.

For the analyses of the stakeholders data of relative humidity, minimum and maximum temperature and global radiation data was prepared. For some of the stakeholders, only data preparation was done (including the choice of possible stations), as they preferred not to give away their own data. For the others, correlations to pollutants and the height gradients of relative humidity were analysed.

In the future, further improvements in homogenisation methods might be needed. At the moment, metadata is not included in ACMANT and there is no possibility to influence the break timing chosen by the software. Moreover, no uncertainty measures are available. A combination of the advantages of ACMANT with the advantages of HOMOP seems desirable. Additionally, analyses of the possibility to use datasets of different parameters homogenised with different methods together would be necessary.

2.2 Contents and results of the project (max. 20 pages)

1) Motivation

To achieve reliable results on the development of the past climate, it is necessary to use well-homogenised data. Homogeneous datasets are influenced by climate alone. Original observational data include effects caused by station location, used instrumentation, the observer, etc. (Aguilar et al. 2003).

To remove those effects, statistical methods are applied to adjust data from earlier periods to the current situation of the station. This process is called homogenisation.

Homogenisation of meteorological data has already been performed for different parameters on monthly resolution since about 20 years. During the last years, the interest in and the activity for homogenised daily data increased (e.g.: COST-Action ES0601 www.homogenisation.org)

In contrast to temperature and precipitation, which have been in the focus of daily homogenisation until now, relative humidity shows a strict upper and lower limit of possible values and the values have limited range. Therefore only small, but nevertheless potentially important, breaks are to be expected, as bigger breaks can be found by data quality control and technical support easily. Moreover, instrumentation of relative humidity measurements is changed more frequently than that of other parameters. Therefore homogenisation methods for relative humidity have to be able to detect small but potentially frequent breaks.

ACRP – Calls for Proposals

2) Objectives of the project

This project aimed to test different methods used for homogenisation of daily data and used within the COST-Action ES0601 on their ability to meet the requirements for homogenising daily relative humidity. If a successful method could be identified, a set of Austrian stations (STARTCLIM-dataset) was to be homogenised. As for those stations where daily temperature extremes and precipitations were already homogenised, this additional homogenised parameter is especially valuable.

In addition, analyses with these data concerning climate change and the effect of the homogenisation itself have been done as well as analyses that were performed in cooperation with stakeholders.

3) Activities performed within the project

a) Surrogate dataset

A surrogate dataset is a dataset of artificially created data that has the statistical properties of real data. The characteristics of the homogeneous data are the statistical probability distribution and the cross- and auto-correlations. Inhomogeneities are added to this data.

▪ *Analysis of real stations*

In order to get information on the size of possible inhomogeneities, parallel measurements for different stations in Austria were analysed to collect information on differences in the distribution of measured values during the different seasons and the year. The stations used are listed in Table 1.

Table 1: List of stations used for analyses of parallel measurements

<i>station</i>	<i>start</i>	<i>End</i>	<i>station</i>	<i>start</i>	<i>End</i>
Kremsmünster	19880101	20081231	Amstetten	19701001	19771130
Salzburg	19920201	20060331	Retz	19940101	19950531
Innsbruck	19920701	20130526	St.Pölten	20030701	20050630
Bregenz	19930101	20130331	Waidhofen	20021001	20041231
Feuerkogel	19891001	20081231	Jauerling	19940101	19950630
Patscherkofel	19930819	20080101	Mallnitz	19870220	20020801
Sonnblick	19881101	20081231	Sillian	19960901	19990930
Kufstein	19920601	20021231	Leibnitz	19941203	20041231

Additionally, the number and temporal distribution of breaks (based on station relocations and changes of instrumentation) was analysed by the study of metadata for the stations distributed over Austria.

Moreover, 6 networks of 4 to 6 highly correlated Austrian stations were created, which represent different regions of Austria (Figure 1). For those networks, a homogeneity test was executed on the monthly data with HOMER (Mestre et al., 2013), created within the COST-Action ES0601. This data was used in the following to create surrogate datasets with 100 years of length in different complexities by the project partner Victor Venema at the University of Bonn.

ACRP – Calls for Proposals




Figure 1: Networks used for Surrogate dataset

■ *Creation of the surrogate datasets*

A stochastic simulation method (IAAFT, iterative amplitude adjusted Fourier transform (Schreiber und Schmitz 1996, Schreiber and Schmitz 2000, Venema et al 2006a-c)) was used to create the long-term homogeneous surrogate dataset from the short homogeneous station data of the different networks. The method allows simulating the exact distribution of the data, which is especially important for relative humidity, where the maximal value of 100% must not be exceeded.

The autocorrelations are constrained by reproducing the power spectrum. The empirical distribution of the measurement is used as input to the IAAFT algorithm. The seasonal cycle is modelled by simulating an empirical distribution for every month. The distribution of the difference time series (cross-correlations) also has a seasonal cycle this way and together with the power spectrum models the cross-correlation matrix.

Judging from the way the spectrum is computed from short time series, variability on decadal scales is likely too weak. Therefore we explicitly model the decadal variability as a smooth noise time series with a power law power spectrum with exponent -4. This time series is normalised to a mean of zero and its standard deviation is about 6% of the variability of the homogeneous data. The same variability was added to every time series in one network.

The statistical properties of the century long surrogate datasets accurately match those of the stations they are based on. This means that the autocorrelation function and the distribution match well and only the cross correlations between the stations are too low over the entire seasonal cycle. This indicates that the convergence of the iterative algorithm was not ideal, which is likely due to the strongly non-Gaussian distribution of the humidity data. The deviations seem acceptably small for this application, however. The cross correlations of the surrogate data could represent a network with a slightly different geographical configuration. Nevertheless, higher correlations can also be found for some networks and months.

■ *Including of inhomogeneities*

Three different kinds of inhomogeneities have been created: “deterministic”, “stochastic” and “realistic”.

ACRP – Calls for Proposals

In the realistic case, three kinds of missing data are included: 1) The number of stations increasing with time. 2) During the Second World War (1940-1946) more data are missing than during other periods. In 1946 75% of stations have missing data and stations with missing data in one year, have a 75% probability of having missing data in the preceding year as well. 3) Random missing data was inserted. The probability of one new missing data value being inserted is 0.0016. The next day has a 75% probability of also being missing.

Inhomogeneities are modelled as a power law power spectrum with exponent -4; see x-axis of Figure 2. In addition they have a seasonal cycle, which is generated by repeatedly smoothing white noise (periodic boundary conditions) and normalising it; y-axis of Figure 2. After adding the perturbations (example see Figure 2), tapering is applied to avoid relative humidity values above 100%.

In the case of the stochastic and realistic dataset, white noise was added to the inhomogeneities. The standard deviation of this white noise was modelled in the same way as the deterministic inhomogeneities.




Figure 2: Example of break as function of homogeneous value and day of the year

In the first two datasets, breaks are inserted with a frequency of 1 break/13years. In the realistic dataset the breaks are more frequent in the beginning of the time series.

The mean of the break size changes linearly in time to create a bias in the relative humidity of 2% over the century. To reduce problems with the upper level, it was assumed that old values were biased 2% too low. Information on real long term biases is not available, but including a bias was seen as desirable to be able to assess how well the methods would handle a trend if there was one.

b) Handling of simultaneous breaks

At the turn of the year 1970 to 1971, the evening observation in the whole meteorological network of Austria was changed from 9pm to 7pm. As this was done simultaneously, no relative homogenisation can be applied. Therefore an alternative method for adjusting these breaks was developed using hourly data for training and validation. To adjust the 9pm value, different methods like spline interpolation, similarity approach (Brandsma and Können, 2006), linear interpolation between the adjacent measurements, multiple linear regression model and seasonal multiple linear regression model were tested. The best results were achieved using

ACRP – Calls for Proposals

the seasonal multiple regression model, using the measurements of this day and the 7am measurement of the following day as well as the time of sunrise and sunset. The original range of RMSE for those stations was reduced from -0.5% to 3% to the range of -0.5% to 0.5%.

Nevertheless, the coefficients seem to depend on small scale features, so that no general coefficients could be found. Therefore only those stations for which hourly values are available before and after 1971 could be corrected. Only 17 stations have been corrected in this manner. The corrections vary between $\pm 11\%$, with one third of the corrected data points needing no correction.

c) Homogenisation

▪ *Methods*

The following methods, being used in the comparisons of the COST-Action ES0601 on daily temperature data, have been applied to the surrogate dataset of relative humidity (daily average):

HOMOP: (Gruber et al., 2009, Nemec et al., 2012)

Break detection is done with PRODIGE (Caussinus & Mestre, 2004), applying a penalised likelihood method comparing the candidate stations with as many reference series as possible. Three different penalty-terms are used. And in this project two different criteria were applied to decide if a break signal is strong enough to indicate a break. The usual criterion for a break is that the signal is shown by at least half of the reference stations in at least two seasons by at least 2 penalty term versions. With the second criterion used in this project, only the results of one penalty term has to indicate a break in two seasons for more than half of the reference stations.

Two different break correction methods have been applied: In one case, the correction was done by SPLIDHOM (Mestre et al, 2011) using a spline regression, resulting in different corrections for different values. Outside of the data range of the reference period, a constant correction was applied.

The second correction method is INTERP (Vincent et al. 2002), calculating daily adjustments by interpolation of monthly adjustments.

Moreover, two different options of reference periods have been tested: in one case the maximum possible reference length was used, and in the other case the reference period was reduced to a maximum of 5 years to minimize the possibility of breaks in the reference series within this period.

ACMANT: (Domonkos, 2011)

This fully automatic method was developed for temperature data in mid- and higher latitudes. Composite data of at least 30years length are used as reference. The detection is based on PRODIGE (Caussinus & Mestre, 2004), but there are some differences:

ACRP – Calls for Proposals

- 1) the reference series are prehomogenised
- 2) a bivariate detection (for annual values and the seasonal amplitude) is used
- 3) homogeneous parts of the time series longer than 3 years are detected separately from inhomogeneities that have a strong signal, but result in a short homogeneous part of the time series (only 3-60month between two breaks)
- 4) a composite reference is used instead of pairwise comparison of stations

The correction is based on ANOVA, excluding breaks that are not significant and correcting only those significant ones. The adjustments are calculated from monthly correction values.

The stations of the whole network are homogenised using this method.

MASH: (Szentimrey, 1999)

This iterative method uses all stations as candidate stations and reference stations. In each iteration, the homogenisation is based on the results of the last step. The homogeneity test is done using test statistics. Daily adjustments are based on monthly ones.

To apply this method it was necessary to divide the measurements by 10.

The method was applied in two ways: In one version a fully automatic script was started, which is not recommended by the authors, and in the second version the homogenisation was done interactively, changing break points if they seemed unrealistic. As no metadata was available for the tests of the benchmark dataset, it seemed possible that the fully automatic process might lead to better results than a human without any additional information on the breaks.

PROCLIM: (Štěpánek, 2008)

This software combines many different detection and correction methods. Following the recommendation of P. Štěpánek, 3 different break detection methods (SNHT, bivariate, t-test) and their combination and 2 correction methods were chosen. One of the corrections used daily adjustments based on monthly corrections and the second one used the q-q-method for directly determining daily adjustments.

A total of 17 different methods (including different options for one and the same algorithm) were applied to the surrogate dataset in order to find the method best suited for homogenisation of relative humidity. In Table 2 the different methods and their abbreviation in the different plots are listed.

Table 2: Abbreviations used in the plots for original data and homogenisation methods

Method	Nr.	Abbreviation
ssearched homogeneous time series	0	soll
Inhomogeneous original time series	1	Ausgang
ACMANT	2	ACMANT
MASH	3	MASH
MASH with interaction	4	MASHeingriff
HOMOP using Splidhom	5	HOMOP
HOMOP using Splidhom with changed break detection criteria	6	HOMOP_get1crit, HOMOP_crit2

ACRP – Calls for Proposals

HOMOP using Splidhom with shortened reference period	7	HOMOP_get1crit_kref, HOMOP_crit2_kurzRef
HOMOP using Vincent	8	HOMOPvin
HOMOP using Vincent with changed break detection criteria	9	HOMOPvin_get1crit, HOMOPvin_crit2
HOMOP using Vincent with shortened reference period	10	HOMOPvin_get1crit_kref, HOMOPvin_crit2_kurzRef
PROCLIM with merged break detection and monthly correction	11	PROCLIM_m102
PROCLIM with SNHT break detection and monthly correction	12	PROCLIM_a102
PROCLIM with bivariate break detection and monthly correction	13	PROCLIM_b102
PROCLIM with t-Test break detection and monthly correction	14	PROCLIM_t102
PROCLIM with merged break detection and qq correction	15	PROCLIM_m106
PROCLIM with SNHT break detection and qq correction	16	PROCLIM_a106
PROCLIM with bivariate break detection and qq correction	17	PROCLIM_b106
PROCLIM with t-Test break detection and qq correction	18	PROCLIM_t106

▪ Testing homogenisation methods

The methods are validated using the surrogate datasets. They have been created artificially, but represent the statistical characteristics of the real data. In contrast to real data, the breaks are known and in comparing the inhomogeneous surrogate data and the homogenised data to the homogeneous surrogate data (the surrogate data, before breaks have been included), the effect of the homogenisation can be measured.

Two abilities of the homogenisation methods have been tested:

- *) break detection
- *) quality of the homogenised dataset (RMSE, trends, variability)

As it is difficult and partly impossible to include the breaks not found by the method itself (e.g. by fully automated systems without any possibility of interaction), no comparisons were done testing the correction methods by using the correct timing of the breaks.

The homogenisation of the surrogate data was done without knowledge about the timing of the breaks.

Break detection

In Figure 3, an example of the break detection for a time series in the deterministic surrogate dataset is shown. In the upper part of the figure, the inhomogeneities of the different breaks of this time series is shown as a function of the measured value and for the different seasons. The timing of the breaks can be found on the x-axis. The multicolour circles in the middle part of the plot show which method detected this specific break and in the lower part all the breaks detected by the different methods are displayed (colours following the same code as the multicolour cycles). A break is defined as detected as soon as the detection takes place in the according year or +/-1year. The detection, e.g. in MASH, is done for each month separately, therefore the number of detected breaks is higher than e.g. for HOMOP. This was taken into account by only using one break per year for MASH. Breaks in neighbouring years were not counted as more than one if they could be attached to one and the same break of the surrogate dataset.

ACRP – Calls for Proposals

The ability in detection was calculated using the detection efficiency (Domonkos et al., 2011). The efficiency is defined as the ratio of correctly and falsely detected breaks to the number of searched (inserted) breaks:

$$E_D = \frac{\#breaks_{correct} - \#breaks_{false}}{\#breaks_{inserted}}$$

In the case that all breaks are correctly detected and no additional breaks are detected, E_D is 1. For those methods that detect the breaks for each month separately (PROCLIM, MASH), the E_D -values are strongly negative (Figure 4). A distinct improvement can be noted between MASH and MASHeingriff. Nevertheless, the efficiency stays negative for most of the time series. There are differences between the different detection methods of PROCLIM. In contrast to the other methods, the bivariate method shows a positive efficiency for most of the time series.

Looking at differences in the different networks, it can be noted that the worst results are reached for network 5 and the best for network 4 and 8 in the deterministic case. Regardless, network 5 shows the best, but still negative results, for 2 methods. MASHeingriff improves all the networks in comparison to MASH.

In accordance with expectations, the detection efficiency for realistic networks is less than for the deterministic ones, as the detection is rendered more difficult by missing values, additional noise and different length of the time series.

In the deterministic and the realistic case the methods ACMANT and HOMOP show the highest detection efficiency.

All the methods classified some of the series as homogeneous. Most time series wrongly described as homogeneous occurred with HOMOP taking into account the deterministic and the realistic networks. Taking a look at the deterministic networks alone, MASHeingriff classified more time series wrongly

Additionally, an analysis using ACMANT, MASH (automatic version only) and HOMOP was performed on the homogeneous time series, to see if those would be defined as homogeneous by the methods. All of the methods detected breaks, but their number was roughly an order smaller than in the inhomogeneous case.

Break characteristic

A further analysis was done in order to gain understanding of which characteristics the breaks found by the different methods would have to meet. The size of the break signal, the correlation within the network and the clustering of the breaks within the time series itself and in combination with breaks of the reference stations were taken into account. In order to keep this idea as simple as possible, only the deterministic validation datasets were used. The break was classified as “clustered” if more than one break lay within a time window of 5 years.

ACRP – Calls for Proposals




Figure 3: Examples of break detection of different methods: line plots in the upper part: structure of the induced break for different seasons (legend above) for the different breaks in the year (x-axis), in which the centre of the plot is plotted. The legend for the axis of the plots can be found on the right of the vertical line in 2000). Middle part: Colours of this pie chart show which of the methods have detected a break in this year (± 1 year). The height of the centre of the circles indicates the yearly mean break size (see y-axis for the value) for the year. The orange letters give the mean break size for the seasons (F...spring, S...summer, H...autumn, W...winter). The legend for the pie chart is given on the lower right site of the vertical line in 2000.

Abbreviations for the Methods: A...ACMANT, H...HOMOP, H1...HOMOP with changed break criteria, M...MASH, Me...MASH with interaction, PS....PROCLIM , Pb....PROCLIM bivariate, Pt...PROCLIM t-Test, Pm...PROCLIM merged). The lines of dots in the lower part give the information about all detected breaks by the methods (colours are the same as in the pie chart, diamonds indicate the 4 PROCLIM versions).

Only 4 breaks in the surrogate dataset were not clustered with breaks in the reference time series and those occurred at the beginning of the time series (between 1901 and 1909). Half of them were not detected by any method, the other half was detected by 3-4 methods. About $\frac{3}{4}$ of all the breaks were additionally clustered with breaks of the time series itself.

The percentage of correctly detected breaks increases with the correlation within the network. Moreover, the ratio of correctly detected breaks increases with the absolute value of the ratio of break size to the variance of the time series. In the case of the surrogate dataset of relative humidity, most of the breaks have a signal about the same size as the variance.

ACRP – Calls for Proposals




Figure 4: Detection efficiency for all time series for the different detection methods in the deterministic cases

Break correction

Not all methods have been able to provide complete solutions for all the networks, as for some methods and stations the correlation of the reference series was not sufficient. In the realistic case, the final quality control by MASH had to be skipped as the number of reference stations was not sufficient in these cases. The number of not solvable stations stays nearly the same in the deterministic and realistic case. Most problems had ACMANT and PROCLIM. However PROCLIM was not used in the realistic case for break correction as it gave the worst results in the deterministic case and moreover did not work satisfactorily on all computers.

The improvement by homogenization of RMSE, variance (time series with removed annual cycle were used in this case) and trend of the homogenized time series were analysed. For RMSE and variance, only the part of the time series in which differences should/did occur was taken into account. Therefore the data between the newest break point (first detected or introduced in the surrogate dataset) and the beginning of the time series (oldest data point) was used. This causes slight differences in the length of the analysed time series with the different methods, but as only means are compared these differences seem of no consequence. The advantage of this method is that long homogeneous parts at the end of the time series (most current data) do not influence the results. All time series for which a homogenization result was achieved were taken into account, including time series classified as homogeneous by the homogenization method.

Trends in relative humidity are (if existing) very small and not significant, but as the improvement of trends is one of the declared goals of homogenization, this parameter was also looked at.

Following Domonkos et al. 2011 the RMSE-efficiency and the trend-efficiency were calculated:

$$RMSE_{Efficiency} = \frac{RMSE_{original} - RMSE_{homogenised}}{RMSE_{original}} * 100$$

ACRP – Calls for Proposals

$$Trend_{Efficiency} = \frac{(\Delta Trend_{original}) - (\Delta Trend_{homogenised})}{(\Delta Trend_{original})} * 100$$

Only ACMANT could solve the bigger part of the stations with a positive RMSE-efficiency (Figure 5) in the deterministic cases. The ratio between positive and negative efficiencies is similar between HOMOP and MASH. PROCLIM gives the worst values of RMSE-efficiency and only a small portion of the stations can be improved.




Figure 5: RMSE-Efficiency for all time series in the deterministic networks for the different homogenisation methods. Green shows the number of time series with negative and positive efficiency. Time series with no change in the RMSE are not included there.

In the realistic case ACMANT results in the highest number positive influenced time series (about half of the time series). All other methods did not improve a single time series.

In the case of Trend-efficiency, all the methods improved about 1/2 to 2/3 of the time series. The best results are reached by ACMANT and HOMOP using spline corrections with the complete reference period. The highest number of negatively influenced time series resulted from PROCLIM using monthly correction with the breaks from the break detection methods t-test- and merged-break detection. In the realistic case, MASH improves most time series (improved: 116/worsened: 44). The ratio of ACMANT is comparable, but less stations have been

ACRP – Calls for Proposals

homogenised by this method (improved: 91/worsened: 43). HOMOP shows the worst results with only about half the stations improved.

A comparison of the differences of the variance between the optimal solution and the homogenised time series resulted in the deterministic cases in nearly no changes compared with the inhomogeneous data for MASH, MASHeingriff and HOMOP with Vincent correction, short reference period and the lighter break definition. All the other methods show a broader distribution with more extreme differences in the variance of the optimal solution.

In the realistic case (Figure 6), it is still MASH that shows the smallest median of differences, but MASH is also worsening some of the time series. Once again, the human influence made MASH results slightly worse. When using ACMANT, the median of the homogenised series is similar to the median in the inhomogeneous cases, but some outliers show distinctly higher differences. The HOMOP methods give the worst results, showing an increase of variance in most of the time series.




Figure 6:
Differences of the variance compared to the optimal solution for all networks and stations for the realistic datasets. Numbers on the x-Axis give the homogenisation methods (see Table 2)

d) Homogenisation of Austrian time series

As can be seen in the results of the tests with surrogate data, only MASH and ACMANT show an ability for homogenising daily averages of relative humidity. Therefore those methods were tried for homogenising real station data.

Networks of 4 stations were built, of which all had a correlation of at least 0.6 and were located within a radius of 200km horizontal and 200m vertical. The period for homogenisation was chosen according to the already available data, starting in 1948 when possible. The period had to be shortened for some stations.

MASH and ACMANT were able to provide solutions for those networks. The choice of the final homogenisation method used in this project was therefore taken somewhat subjectively, as MASH detected the first breaks right at the beginning of the time series, which resulted in very

ACRP – Calls for Proposals

short reference periods. Moreover, the test showed that ACMANT improves the RMSE better than MASH and gives similar results in the improvement of the trends. As relative humidity is often used in combination with other parameters, the improvement of RMSE seems quite important. *Thus, although MASH has the advantage of including information on metadata and therefore of possible break points, ACMANT was chosen.*

Table 3: Information on the homogenised stations and the homogenisation result: For breaks written in orange colour, meta-information within +/-1year can be identified (in case of Hohenau, no long term metadata was available). As valid metadata changes in the humidity instrumentation, changes in the observer and relocations (or the documented construction of building in the surrounding) were used. Station numbers written in blue mark those stations for which a correction of the change in observation time has been done. The subjectively determined quality of the homogenisation is given in colour code (green: ok, orange: ~, red: bad). In the column “#stat” the number of stations in the network is shown.

Station Nr.	Station (lon,lat,height[m])	# stat	quality & comment	# breaks
90500 (1948-2013)	Retz (15°57',48°46',320)	15		6 (1977,1990,1992,1994,1997,2000)
140100 (1960-2013)	Kollerschlag (13°50',48°36',714)	14		9 (1967,1973,1979,1981,1988,1991,2001,2002)
160100 (1948-2013)	Freistadt (14°30',48°30',539)	11		7 (1953,1958,1971,1989,1991,1998,2001)
240000 (1956-2013)	Laa/Thaya (16°23',48°44',184)	19		11 (1963,1967,1968,1979,1983,1986,1991,1993,1997,2003,2009)
260000 (1949-2013)	Hohenau (16°54',48°37',154)	16		6 (1949,1952,1959,1962,1969,2007)
290000 (1949-2013)	Reichersberg (13°23',48°20',351)	12		12 (1950,1955,1958,1962,1965,1976,1979,1985,1988,1995,2000,2004)
341000 (1948-2013)	Pabneukirchen (14°40',48°21',554)	14		6 (1954,1957,1976,1982,1989,2004)
380500 (1948-2013)	Krems (15°37',48°25',203)	22		2 (1973,1995)
501200 (1948-2013)	Kremsmünster (14°08',48°03',382)	17		8 (1954,1962,1989,1992,1996,2001,2004,2008)
590400 (1948-2013)	Wien Hohe Warte (16°21',48°15',198)	16		5 (1962,1970,1987,1991,1993)
599000 (1948-2013)	Schwechat (16°34',48°07',183)	13		4 (1971,1983,1994,2011)

ACRP – Calls for Proposals

630000 (1948-2013)	Salzburg (13°47'48",430)	18		9 (1954,1962,1968,1971,1979,1987, 1994,1998,2010)
691000 (1953-2000)	Großraming (14°31',47°54',379)	8	few ref. in early years	6 (1956,1968,1973,1983,1985,1997)
770400 (1976-2013)	Eisenstadt (16°32',47°52',184)	15		4 (1985,1988,1996,2003)
964000 (1963-2013)	Bad Aussee (13°46',47°37',743)	12		4 (1975,1979,1983,1995)
1180300 (1948-2013)	Universität Innsbruck (11°23',47°16',577)	9	few ref. in early years	2 (1971,1979)
1190100 (1956-2013)	Jenbach (11°45',47°23',530)	14		11 (1962,1968,1970,1976,1977,1982, 1986,1997,2003,2009,2010)
1232200 (1948-2013)	Zell/See (12°48',47°20',751)	17	few ref. in early years	7 (1968,1975,1985,1989,1996,2001,2005)
1500100 (1948-2013)	Mayrhofen (11°51',47°10',640)	11		8 (1955,1964,1967,1976,1980,1992,2004)
1540200	Rauris (13°,47°13',941)	16	few ref. in early years	8 (1961,1964,1977,1984,1989,1993,1999, 2003)
1571200 (1984-2013)	Tamsweg (11°35',47°08',1025)	7		3 (1997,2008,2010)
1591000 (1952-2013)	Stolzalpe (14°11',47°07',1291)	8	few ref. in early years	3 (1980,1997,1999)
1630000 (1961-2007)	Lobming (15°11',47°03',414)	9		4 (1975,1985,1994,1999)
1640000 (1965-2013)	Flughafen Graz (15°26',47°,340)	13		4 (1971,1990,1997,2000)
1641200 (1965-2013)	Universität Graz (15°27',47°05',367)	12		10 (1968,1973,1978,1980,1983,1986,1992, 1996,1999,2003)
1650000 (1961-2013)	Gleisdorf (15°42',47°07',377)	14		5 (1977,1984,1992,1998,2005)
1671100 (1955-2013)	Wörterberg (16°06',47°14',404)	19		7 (1963,1973,1981,1986,1997,2003,2008)
1770000 (1948-2013)	St. Jakob/Defreggental (12°21',46°55',1383)	5	too few ref.	2 (1996,1997)
1790100 (1948-2013)	Lienz (12°48',46°50',661)	13	few to no ref. in early years	7 (1965,1973,1986,1990,1995,1998,2004)
1810000 (1955-2001)	Kolbnitz (13°19',46°52',603)	12		4 (1957,1963,1967,1988)

ACRP – Calls for Proposals

1880500 (1951-2013)	Preitenegg (14°55',46°56',1034)	8	few ref. in early years	7 (1969,1993,1998,2001,2004,2007,2009)
1980000 (1961-2013)	Reisach (13°09',46°39',646)	9	few ref. in current years	5 (1971,1974,1986,1988,1995)
2021200 (1964-2013)	Klagenfurt (14°19',46°39',450)	16		7 (1967,1975,1976,1998,2006,2008,2010)
2040000 (1961-2013)	St. Michael ob Bleiberg (14°46',46°34',532)	19		5 (1980,1986,2007,2009,2011)

The networks were expanded with further reference stations to improve the homogenisation results.

In Table 3, the stations for which a homogenisation was possible are displayed along with their coordinates, the number of reference stations used (not all of them were available through the whole period) and some information on the homogenisation and the data series.

Some stations were excluded as the available time series was too short, too incomplete or because a sufficient number of reference stations was not found. An examples of the homogenisation is displayed in Figure 7.

The quality of the homogenisation has not been assessed by the software so far. Therefore all time- series are to be called homogenised; nevertheless the quality of the homogenisation was subjectively determined by the length and number of available reference series.

The change in the observation time in 1971 may still be found in some of the stations. This is especially true for those stations that have been corrected for this change. As ACMANT doesn't offer the possibility to influence the used break points, the influence of the not corrected stations cannot be counteracted.

e) Analyses of relative humidity

▪ Stakeholder

In cooperation with 4 stakeholders analyses of relative humidity showing the large potential of humidity series were defined.

Where possible and necessary, homogenised temperature values of the project HOM-START (www.zamg.ac.at/cms/de/forschung/klima/datensaetze/homstart; Nemec et al. 2012) have been used and were prolonged by original data for the recent years. Daily mean temperature was calculated by averaging T_{\min} and T_{\max} . For relative humidity in the original data, time correction was applied where possible and necessary.

The annual cycle (on daily base) and trend has been removed for all time-series (if not stated otherwise below) for correlations. Both actions have been taken to remove "wrong" correlation information as all parameters are influenced by the annual cycle and trends. Without removing those effects, the correlation would be higher than by looking at the signal of the parameter itself (Table 5). As not all the parameters are normally distributed, the spearman's rank correlation has been used. The statistical significance is given by the p-value.

ACRP – Calls for Proposals




Figure 7: Example of a real homogenised time series (Lobming). Upper panel: original data (Lobming: red) and reference series (black); middle panel: as upper panel but homogenised data; lower panel: Lobming original data (black) and homogenised (red). Displayed annual means.

The analyses were repeated with the homogenised data if necessary, but no essential changes occurred.

The first cooperation took place with Prof.DI.Dr. Hans-Peter Hutter of the Medical University of Vienna. He was interested in the influence of relative humidity and temperature on the concentration of air pollutants NO₂ and PM₁₀ in urban regions. The cities of Graz, Linz and Vienna were chosen. Further investigation concerning physical health indicators like the number of discharges from hospital and mortality (cardiovascular and respiratory) will be done by the stakeholder himself if necessary.

The data on NO₂ and PM₁₀ was provided by the Federal government of Styria, Upper Austria and the Conservation department of the Vienna city administration.

Information on NO₂ and PM₁₀ concentrations is not available at the same locations as the meteorological data. The locations of the stations used are given in Table 4.

city	meteorological	NO ₂	PM ₁₀
Graz	Graz airport	Graz/Sued Tiergartenweg	
Linz	Hörsching	Traun/Tischlerstraße	
Vienna	Hohe Warte	Hohe Warte	Wien/Schafbergbad

Table 4: Location of stations used for investigation of pollutants and meteorological parameters

The correlations between the meteorological parameters and the pollutants have been calculated (Table 5) for the time period for which the data of the pollutant was available.

Differences in the correlation can be seen according to seasons. Additionally, a delay can be found for some of the time series, resulting in higher correlations when calculating the correlation between the pollutant and the meteorological value of the following day. The influence of season and shift differs for the different locations.

ACRP – Calls for Proposals

Table 5: Spearman correlation between the time series of relative humidity/temperature and air pollutants (first value in column “correlation” and “significance” for original data, second value for time series with removed annual cycle and trend; statistically not significant correlations are marked with an “-“)

city (pollutant)	Investigation period	Correlation (RH)	Significance (RH)	Correlation (T)	Significance (T)
Graz (NO ₂)	20030425 - 20140531	-0.60/-0,01	0,1%/-	0.31/-0,05	0,1%/0,5%
Vienna (NO ₂)	19880101 - 20140531	-0.30/0,07	0,1%/ 0,1%	0.22/0,07	0,1%/ 0,1%
Linz (NO ₂)	19900101 - 20140531	-0.44/0,01	0,1%/-	0.24/-0,05	0,1%/ 0,1%
Graz (PM ₁₀)	20030425 - 20140531	-0.51/0,07	0,1%/ 0,1%	0.29/-0,04	0,1%/5%
Vienna (PM ₁₀)	20020412 - 20140531	-0.26/-0,00	0,1%/-	0.15/0,08	0,1%/ 0,1%
Linz (PM ₁₀)	20001212 - 20140531	-0.20/0,04	0,1%/0,5%	0.01/-0,13	-/ 0,1%

The correlations between meteorological data and concentration of air pollutants do not include quite important processes that influence the air pollutants. In the case of PM₁₀, the main influencing factors are distant sources of air pollutants and their advections by large-scale atmospheric conditions, or also neighbouring sources like construction sites or winter road gritting. Moreover, in this analysis no distinction between potential periods with and without temperature inversions was drawn.

In the case of NO₂, chemical reactions have an influence on the concentration of the pollutant as well as local influencing factors (e.g. road traffic, which might explain the period of small concentration around Christmas holidays)

The second analysis was done for Dr. Meinhard Breiling of the Technical University of Vienna. His research is focused on the artificial snow production and therefore his interest lies in the climatological information on the conditions to produce snow. In this project, an analysis on monthly and seasonal (winter half-year: October to March) basis of vertical gradients of air temperature and relative humidity between stations pairs (Table 6) was done.

To reduce the possibility of breaks in the high level station, for which no homogenised data is available, a common period (19970101-20081231) was chosen for all stations for which no relocations took place.

Low level station	Height [m]	High level station	Height [m]
Graz university	366	Schöckl	1436
Innsbruck university	578	Patscherkofel	2251
Rauris	934	Sonnblick	3109

Table 6: Information on stations used for vertical gradients

Looking at the differences between high and low level stations, the medians for the months and season are, with the exception of January, negative for Rauris and Sonnblick, meaning that relative humidity is predominantly higher at high elevations. The lowest medians are exhibited in February and March.

ACRP – Calls for Proposals

For temperature, as expected, values at higher stations are lower than at lower stations. Negative values occur only seldom and are small. As only two stations have been used for these analyses, no information about the height of inversions was available. As radio sounding data cannot be used easily as a comparison for daily means, no further attempts in this direction were made. The station pair Graz/Schöckl shows the highest percentage of positive temperature gradients (10% of the days). The height between these stations seems therefore to be not too far from the height of usual temperature inversions, as the effect was not counterbalanced by the usual temperature reduction by height. The most/strongest temperature inversions occur in December and January.

As only the winter half year was analysed, no correction for annual cycle and trends have been done when calculating the correlations (Spearman) between temperature and relative humidity gradients.

For the other two stakeholders (Dr. Thomas Bednar of the Technical University of Vienna, and Dr. Thomas Cech of the Federal forest research centre (BFW)), only data preparation was necessary, as both of them wanted to use the data not available for the public within their own software.

In the case of Dr. Bednar, the topic of the investigation was the influence of daily T_{\min} and T_{\max} , relative humidity and global radiation on construction materials. The locations and periods of data provided are listed in Table 7.

The topic of Dr. Cech is the dieback of ash trees in Austria and therefore correlation with relative humidity is of importance. The investigations focus on Lower Austria, as monitoring data of the dieback is available there for 2000 to 2006. Moreover, regional gradients from west to east should be analysed.

Table 7: Stations and periods of data provided for Dr. Bednar

station	Tmin, Tmax	RH	radiation
Vienna -Hohe Warte	19480101-20091231	18720101- 20141231	19600101-20141231
Klagenfurt	19480101-20091231	19500401- 20141231	19690101-20141231
Gmunden	19300101-20141231	19480101-20141231	19700101-20120214
Feuerkogel	19300119-20141231	19300118-20141231	19700101-20141231

Stations were chosen that were located as closely as possible to the BFW monitoring areas in a similar elevation distributed across Lower Austria from west to east. The chosen stations are listed in Table 8. All available data, starting at least in 1948 and reaching until 2014, was used.

Table 8: Stations chosen for Dr. Cech

station	Hohenau	Krems	Kremsmünster	Schwechat	Amstetten
Monitoring site	Hohenau	Rossatz	Haag	Göttlesbrunn	Winklarn

ACRP – Calls for Proposals

▪ Additional analyses

The trend is influenced by the homogenisation of all of the stations (Figure 8). All trends in this figure are calculated using all available data for the stations, therefore the differences in the trends are influenced by the time span that was available.

Before homogenisation, more than 10 stations showed a positive trend, and after homogenisation only three stations (1500100, 177000 and 964000) show a positive trend signal. 964000 is the only station for which the trend changed from a negative sign to a positive one by homogenisation. For all the other stations the trend became more negative. Nevertheless the slope is less than 0.2% per year. The biggest change took place for 1571200 (~0,3% per year). All trends are statistically significant in the homogenised dataset. Using a 95%-confidence interval the trend change is only statistically significant for 25 stations.

Looking at the different seasons, the most positive trends occur in autumn and winter while nearly no positive trend can be found in spring and summer. The change to a positive trend for 9640000 can be found in each season. For 4 other stations changes to positive sign can only be found in one to two seasons.

Taking into account only those stations for which the time series starts in 1948, the homogenised and original trend can be seen in Figure 9. In the beginning of this homogenised time series an increasing or constant level of relative humidity can be found. Between 1970 and 1990 a decrease can be observed, afterwards a relatively constant value is reached again. The original time series start with a lower value of relative humidity. Therefore the decrease is smaller after 1970, ending in a slight increase after 1980 and a constant value in the present.




Figure 8: Information on annual trends of original and homogenised dataset. Station numbers given on the x-axis, the slope [% per year] on the y-axis. Black and grey points give original values, red and orange show the homogenised data. Grey and orange crossed circles give the 95%-confidence interval. The values give the original trend values.

The annual cycle of relative humidity shows only slight changes (Figure 10). For all months, the median shows higher relative humidity in the homogenised data than in the original data. The difference is larger in the warmer months (May-July |median of differences|>3%) than in the cooler (Dec-Feb |medians of differences|<=2%) ones.

ACRP – Calls for Proposals

No height dependence seems to occur in the median and variability of relative humidity in the different months.




Figure 9: Annual time series for 5 stations (given on the lower part of the figure) starting 1948 and trends. Slopes [% per year] and p-values for the slopes are given at the lower part of the figure.




Figure 10: Annual cycle of relative humidity values for original (left) and homogenised (right) data

A comparison between the monthly RH-data homogenised within the framework of HISTALP in 1997 and the daily homogenised data from this project (averaged over the single months for equal resolution) was done for all 4 stations that are available in both datasets (Innsbruck University, Kremsmünster, Wien Hohe Warte, Graz University). The differences between the original data (CC-IMPATY minus HISTALP) have values between -7% and 5% (mean: -3%) due to different calculation of monthly means in both datasets (using daily means in CC-IMPATY and all observation-time measurements of the months for the HISTALP-data). Comparing the homogenised time series from the beginning of the available series until 2013 (using unhomogenised data before 1997 for HISTALP) showed that the differences have a range of -15% and 14% (Table 9)(mean: -4%). In most months, mean monthly humidity is higher in

ACRP – Calls for Proposals

HISTALP than in CC-IMPATY. Only in Kremsmünster did roughly half the months show means indicating that relative humidity is lower in HISTALP. No common seasonal cycle can be seen between the differences of HISTALP and CC-IMPATY in the different stations. When looking at trends for the single months, they are very similar for three of the stations having mean absolute errors of 0.05%/y. Only Graz University has differences of $\geq 0.1\%/y$ for each month.

	Kremsmünster	Wien Hohe Warte	Innsbruck Uni	Graz Uni
M1	-6/-2/2	-5/-1/1	-8/-4/-1	-13/-6/-1
M2	-5/-3/1	-12/-5/0	-9/-5/-2	-13/-7/-2
M3	-6/-3/-1	-10/-5/-2	-11/-6/-4	-13/-8/-3
M4	-10/-3/5	-15/-10/-2	-12/-6/-2	-14/-9/-4
M5	-4/5/10	-10/-6/-2	-12/-5/0	-12/-5/-1
M6	-4/6/14	-9/-3/-1	-11/-5/-2	-13/-5/0
M7	-5/2/10	-10/-5/-2	-11/-5/1	-14/-5/-1
M8	-8/-3/5	-10/-5/-2	-11/-5/-1	-14/-7/-1
M9	-5/0/5	-5/-2/6	-9/-5/4	-12/-5/-1
M10	-4/0/5	-6/-2/3	-8/-4/-2	-13/-7/-2
M11	-5/-3/2	-9/-2/3	-7/-3/-1	-14/-6/-2
M12	-4/-1/4	-5/-1/2	-7/-5/-2	-11/-5/-1

Table 9: Distribution of differences (CC-IMPATY minus HISTALP, min/mean/max [%]) for each month (M1-M12) for homogenised monthly data

2.3 Conclusions to be drawn from project results (max. 5 pages)

Findings:

Homogenisation methods:

- *) The tests using surrogate data showed that in break detection ACMANT and HOMOP performed best with fewer time series wrongly classified as homogeneous by ACMANT.
- *) The number of breaks was reduced significantly by human interaction with MASH in comparison to the automated version. That led to most wrongly as homogeneous classified time series in the deterministic case. In the realistic case the number of wrongly classified time series was similar to the automatic version.
- *) MASH and PROCLIM detect more breaks than ACMANT and HOMOP, although their characteristic to detect the breaks in each month separately was taken into account by summing up each break within 1 year as only 1 break.
- *) The location of the network (inner alpine, stations located nearby or farther from each other,...) seems of less importance to the break detection than the characteristic of missing data, noise,...
- *) All methods tested on their ability to recognise homogeneous time series found breaks in the homogeneous, deterministic surrogate dataset. Nevertheless, the number of breaks was significant (1 order) smaller than in the inhomogeneous cases.

ACRP – Calls for Proposals

- *) Almost all breaks in the surrogate networks were located within 5 years of at least one break in the reference series and more than 2/3 of the breaks lay within 5 years of a break in the candidate time series itself.
- *) The probability of detection increases with increasing correlation of the candidate station and the reference stations and the size of the break signal in comparison to the noise.
- *) ACMANT was the method that provided solutions for the least number of stations.
- *) The network with the most problems was network 8, with only 4 stations.
- *) Of all the methods, only MASH solved all stations for all networks.
- *) In which networks some stations could not be solved depended on the methods and not on the network.
- *) In the realistic case, the quality control usually done by MASH could not be used due to “too few reference stations”.
- *) The RMSE can be improved in the deterministic case (by ACMANT, MASH and HOMOP in 2/3 to 1/2 of the stations), but ACMANT is able to improve the time series (~1/2 of them) in the realistic case.
- *) The trend is improved by all methods for about 2/3 to 1/2 of the stations. The ability to improve the trend decreases in the realistic case, but still about 1/2 of the stations are positively influenced.
- *) There is nearly no influence in the performance of HOMOP with different correction methods, break detection criteria and length of reference period.
- *) The influence of human interaction in MASH is high. Profound induction training is therefore necessary to use this homogenisation tool and should not be done without knowledge of metadata.

Homogenisation of real data:

- *) Daily relative humidity data of 34 Austrian stations has been homogenised.
- *) About half of the breaks can be explained by metadata.
- *) The positive effect of the correction of the time of observation bias was counter acted by the homogenisation due to the higher number of reference stations without this correction.

Analyses:

- *) There seems to be no or only very little influence of relative humidity on pollutants like NO₂ and PM₁₀.
- *) The homogenisation has an important influence on the trend of the relative humidity in the last years, by making the trend negative in most of the cases.
- *) While the median values of relative humidity in the different months increase, the annual cycle remains similar, with a minimum in April and May and the highest values in winter.
- *) No height dependence seems to exist in the median of original and homogenised relative humidity values.

ACRP – Calls for Proposals

- *) A comparison with monthly HISTALP-data showed that the differences between the two datasets increased by the homogenisation.
- *) Most mean monthly relative humidity values are higher in HISTALP than after the daily homogenisation.
- *) The consistency in trend analyses in individual months is varying. While some month at some station show absolute differences of 0.01%/year, other months at other stations can have differences up to 0.19%/y.

Further steps by the project team:

- *) A comparison between the homogenised data of temperature and relative humidity would be needed to determine if those datasets homogenised with different methods can be used in combination.
- *) Further intercomparison between different homogenisation methods are needed for additional meteorological parameters. And additional methods (e.g. the Swiss method including weather patterns) are of interest.
- *) In case there are additional needs of the stakeholders concerning the analyses done in this project, the project team will of course offer the necessary support.
- *) The results on the trends should be further verified by a comparison to trends resulting from annual or monthly homogenised datasets.

Important points for other target groups:

- *) The tests showed that the methods perform quite differently for the same conditions. The comparison of the methods is of interest to other homogenisation groups in the community. On the one hand, the improvement of methods can be a topic, on the other hand these tests showed that homogenising a new parameter needs profound preparatory work to find the best suited method.
- *) The analyses done with the stakeholders allow them to determine if this topic is fit for further research.
- *) The homogeneous data of relative humidity will be available to the public as soon as published and can therefore be used by everyone who is interested in relative humidity.

ACRP – Calls for Proposals

Bibliography:

- Brandsma T., Können G.P., 2006: Application of nearest-neighbor resampling for homogenizing temperature records on a daily to sub-daily level. *Int. Journal of Climatology*, 26,75-89
- Caussinus H, Mestre O., 2004: Detection and Correction of artificial shifts in climate series, *Journal of the Royal Statistical Society, Series C*, 53,405-425
- Domonkos P., 2011: Adapted Caussinus-Mestre Algorithm for Networks of Temperature Series (ACMANT), *Int. Journal of Geosciences*, 2, 293-309
- Gruber C., Auer I., Böhm R., 2009: Endbericht HOM-OP Austria, Aufbau und Installierung eines Tools zur operationellen Homogenisierung von Klimadaten mit Annexen: Gruber C., Auer I, 2009: Description of a procedure for the homogenization of daily temperature time series, Annex zu Endbericht HOM-OP-Austria; Gruber C., Auer I., 2009: Working instructions for HOMOP-A Procedure for the Homogenization of Daily Temperature Time Series. Annex 2 zu Endbericht HOM-OP Austria; Gruber C., Auer I., 2009: Working instructions for HOMOP – A Procedure for Homogenization of Daily Temperature Time Series: Extensions, Annex3 zu Endbericht HOM-OP Austria
- Mestre O., Gruber C., Prieur C., Caussinus H., Jourdain S., 2011 : SPLIDHOM : A Method for Homogenization of Daily Temperature Observations, *J. Appl. Meteor.Climatol*, 50, 2343-2358
- Mestre O., Domonkos P., Picard F., Auer I., Robin S., Lebarbier E., Böhm R., Aguilar E., Guijarro J., Vertachnik G., Klancar M., Dubuisson B., Stepanek P., 2013: HOMER: a homogenization software – methods and applications, *Quarterly Journal of the Hungarian meteorological Service*, 117,47-67
- Nemec J., Gruber C., Chimani B., Auer I., 2012: Trends in extreme temperature indices in Austria based on a new homogenized dataset, *Int. Journal of Climatology*, doi:10.1002/joc.3532
- Schreiber T., Schmitz A., 1996: Improved surrogate data for nonlinearity tests, *Phys. Rev. Lett.* 77, 635-638
- Schreiber T., Schmitz A., 2000: Surrogate time series, *Physica D*, 142, 346-382
- Štěpánek P., 2008: ProClimDB – software for precessing climatological datasets, CHMI, regional office Brno. [Http://www.climohom.eu/ProcData.html](http://www.climohom.eu/ProcData.html)
- Szentimrey T., 1999: Multiple Analysis of Series for Homogenization (MASH), *Proceedings of the Second Seminar for Homogenization of Surface Climatological Data*, Budapest, Hungar; WMO, WCDMP-No 41, 24-46
- Venema V., Meyer S., Gimeno GS, Kniffka A., Simmer C., Crewell S., Löhnert U, Trautmann U., Macke A., 2006a: Surrogate cloud fields generated with the Iterative Amplitude Adapted Fourier Transform algorithm, *Tellus*, 58A, 104-120
- Venema V., Bachner S, Rust HW, Simmer C., 2006c: Statistical characteristics of surrogate data based on geophysical measurements, *Nonlinear Proc. Geophys.*, 13, 449-466

ACRP – Calls for Proposals

Venema V., Ament F., Simmer C, 2006c: A Stochastic Iterative Amplitude Adjusted Fourier Transform Algorithm with improved accuracy, Nonlinear Proc. Geophys. 13, 247-363

Vincent L.A., Zhang X., Bonsal B.R., Hogg W.D., 2002 : Homogenization of Daily Temperature over Canada, Journal of Climate, 15,1322-1334